

bio west	Technical data sheet	Ref : FT.X0520an Page : 1/2
	Created : H. Kerhervé	Version 01 Date de la version : 21/05/05

**Dulbecco's Phosphate Buffered Saline 10X
w/ Magnesium w/ Calcium Chloride**

CAT N : X0520

Theoretical pH : 3.3 ± 0.3

Osmolarity : >1600 mOsm/l

Colour : clear colorless solution

Storage conditions : +20°C

Shelf life : 48 months

Sterility tests :

- Bacteria in aerobic and anaerobic conditions
- Fungi and yeasts

Endotoxin : < 1 EU/ml (<0.1 ng/ml)

Composition : Displayed on website and in catalog; also available on request

Recommended use :

Use aseptic techniques when handling this buffer.

Product is provided for laboratory use only, and not for drug, human or veterinary use.

Applications :

Dulbecco's Phosphate Buffered Saline (DPBS) is intended for use in the maintenance of mammalian cells where a chemically defined, balanced salt solution provides an environment that will maintain the structural and physiological integrity of cells in vitro. DPBS is commonly used in cell enumerations as a diluent, for rinsing cells and as a buffer in many chromatographic procedures.

biowest	Technical data sheet	Ref : FT.X0520an Page : 2/2
		Version 01

Uses :

Supplements, such as antibiotics, should be added as sterile supplements to the buffer solution. Storage conditions and shelf-life of the supplemented product will be affected by the nature of the supplements.

1. Dilute the 10X buffer 1:10 with cell culture grade water. Water temperature should be 15 to 30°C. Do not heat water. Mix completely.
2. While stirring the solution, adjust the pH 6.9 –7.1 using 1N NaOH. For most cells and culture conditions, the optimal pH of this salt solution after filtration is 7.0-7.4
3. Add deionized or distilled water to the solution to bring it to the final volume. To avoid fluctuations in pH, keep the vessel closed until the solution is filtered.
4. Sterilize the solution using a 0.22 micron filter. A peristaltic pump or an inert gas such as nitrogen can be used to provide positive pressure at 3 – 15 psi. Do not use CO₂ gas.
5. Sterile solutions should be dispensed aseptically into sterile containers. Store buffer solution at 15 to 30°C

Signs of Deterioration:

Buffer solution should be clear and free of particulate and flocculent material.

Do not use if buffer solution is cloudy or contains precipitate.

Other evidence of deterioration may include degradation of physical or performance characteristics.